

When can I stop?

You should keep taking your medication until your doctor tells you to stop.



You should keep taking it, even if you feel better.



If you want to stop taking it, talk to your doctor first.




Important

Some medications do not work well together.

You should always tell your doctor or pharmacist what you are taking.



If your medication looks different it might be a different make.



Alcohol can make you ill if you drink it when you are on medication.



Advice and Support

If you have any questions or worries about your medication talk to...

your doctor




or



the pharmacist at your local chemist

This leaflet was created by Michelle Forster, Betsy Wilkie, Dr Andre Strydom, Caroline Edwards and Dr Ian Hall.

The drawings are by Sally Poole, and from the CHANGE Picture Bank.



These people helped to make the leaflet easier to read: John Stevens, Jackie Ryan, Linda Doherty and Eamon Maher.



For more information contact The Elfrida Society, telephone: 020 7359 7443 fax: 020 7704 1358 Produced February 2001

Your medication is called

risperidone or Risperdal



An atypical antipsychotic

This is not an official manufacturer's patient information leaflet.

Why am I taking this?

It can help if you are anxious or paranoid.



It can help if you are too excited or 'hyper'.




It can stop you from hearing voices or seeing things that are not there.

How will it make me feel?

It should make you feel calm.

It could take a month before you feel better.

What will it look like?



How do I take it?

Swallow your tablet with a glass of water.



If you have a liquid you should measure it very carefully.



Ask your pharmacist if you can mix this medicine with a drink.

When do I take it?


Ask your doctor when to take it.

What if I forget to take it?

Take it as soon as you remember.



If it is nearly time for your next dose, only take one dose.



If you have any problems with taking your medication, you should talk to your doctor straight away.

What about side effects?

You might have some of these side-effects...

feeling drowsy

problems with sleeping

feeling restless

problems with sex



If you are worried, talk to your doctor.

IMPORTANT!

These side-effects are more serious...

feeling faint

a rash

a faster heart beat

feeling stiff

feeling shaky

If you have any of these, you must talk to your doctor straight away!

